Write your name here			
Surname		Other name	s
Edexcel Certificate Edexcel International GCSE	Centre Number		Candidate Number
Chemistry Unit: KCH0/4CH0 Science (Double Av Paper: 1C		/4SC0	
Monday 20 May 2013 – Af Time: 2 hours	ternoon		Paper Reference KCH0/1C 4CH0/1C KSC0/1C 4SC0/1C
You must have: Calculator, ruler			Total Marks

Instructions

- Use black ink or ball-point pen.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided - there may be more space than you need.
- Show all the steps in any calculations and state the units.
- Some questions must be answered with a cross in a box ⋈. If you change your mind about an answer, put a line through the box ₩ and then mark your new answer with a cross \boxtimes .

Information

- The total mark for this paper is 120.
- The marks for each question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶

P41554A

PEARSON

Щ
점
I
S
0
Œ
m
п.
뿌
亡

Helium

0

9

2

က

Group

N

Period

N

3

Hydrogen

-	-		_	Ar Argon 18	_	_		_			+		-				
19	<u>L</u>	Fluori 9	35.	Chloring 17	80	Brown	35	127	-	lodin	8	210	¥	Astat 85			
16	0	Oxygen 8	32	Sulfur 16	62	Selenium	8	128	<u>+</u>	Tellurium	25	210	P0	Polonium 84			
4	z	Nitrogen 7	31	Phosphorus	75	Arsenic	8	122	S	Antimony	10	508	窗	Bismuth 83			
12	ပ	Carbon	28	Silcon 14	73	Germanium	32	119	S	E	8	207	8	Lead 82			
-	8	Boron 5	27	Al Aluminium 13	70	Gallium	31	115	므	Indium	46	204	F	Thallium 81			
					65	Zhc	30	112	8	Cadmium	45	201	무	Mercury 80			
					63.5	CC	53	108	Ag	Silver	47	197	Au	Gold 73			
					59	Z	28	106	Pd	Palladium	46	195	ď	Platinum 78			
					29	Sobalt Soba Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Soba Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Sobalt Soba Soba Soba Soba Soba Soba Soba Soba	27	103	뜐	Rhadium	45	192	_	iridium 77			
					95	e r	56	101	æ	Ruthenium	4	190	ő	Osmlum 76			
					55	Cr Mn	52	66	ر ک	Technetium	3	186	Re	Rhenium 75			
					52	Chromium	24	96	Wo	Molybdenum	42	184	3	Tungsten 74			
					15	Vanadium	23	83	S	Niobium	14	181	Ta	Tantalum 73	1000		
					48	Titanium	22	16	Zr	Zirconium	40	179	Ì	Hafnium 72			
					45	Scandium	21	69	>	Yttrium	33	139	La	Lanthanum 57	227	Ac	Actinium
6	æ	Beryllium 4	24	Mg Magnesium 12	40	Sadiy	8	88	Š	Strontium	88	137	Ba	Barium 56	526	Ra	Radium
7	:	Jthium 3	1	Sodium 11	1	X	19				- 1	133	S	aeslum 55	223	ŭ	Francium

Key

Relative atomic mass Symbol Name Atomic number

2

9

BLANK PAGE

Answer ALL questions.

1 Boron is an element in Group 3 of the Periodic Table.

An atom of boron can be represented as ¹¹₅B

(a) Use numbers from the box to complete the sentences about this atom of boron.

3 5 6 11 16


Each number may be used once, more than once or not at all.

(i) The atomic number of boron is ______5 (1)

(iv) This atom of boron contains 6 neutrons.

of alum	inium.		plete each sentence about an at	0111
	fewer	more	the same number of	
Each wo	ord or phrase may	be used once, mor	e than once or not at all.	
(i) Com	pared to an atom	of boron, an atom	of aluminium has	
***************************************	more		protons.	(1)
(ii) Com	pared to an atom	of boron, an atom	of aluminium has	(-/
	more		neutrons.	(1)
(iii) Com	pared to an atom	of boron, an atom	of aluminium has	
***********	Sam	Q	electrons in its outer shell.	(1)
The elec	tronic configurati	on of aluminium is		(5)
□ A	2.3			(1)
B 2	2.2.3			
	.2.8			

2 A student used this apparatus to make and collect a sample of hydrogen gas.



(a) The reaction in the flask can be shown by this word equation.

metal + acid → salt + hydrogen

(i) The name of the salt formed in the student's experiment is

(1)

- A zinc sulfate
- B zinc sulfide
- C zinc sulfite
- D zinc sulfur
- (ii) The student could have used other metals in this experiment.

Place crosses (☒) in **two** boxes to show the names of two other metals that could be safely used to make hydrogen.

(2)

- A copper
- ☑ B iron
- ☑ C magnesium
- D potassium
- E silver

(b) Describe a test to show that the gas collected is hydrogen.				
Use a barning splint to see if p	op.			
(c) Water is formed when hydrogen combines with oxygen.	***************************************			
Balance the equation for this reaction.	(2)			
$H_2 + \dots + H_2 O$	(1)			
(d) Equation 1 represents a reaction using cobalt(II) chloride that can be used to show a liquid contains water.				
Equation 1 $CoCl_2.2H_2O(s) + 4H_2O(l) \rightarrow CoCl_2.6H_2O(s)$				
In this reaction there is a colour change from blue to pink.				
(i) Which word describes both cobalt compounds in equation 1?				
☐ A anhydrous	(1)			
■ B aqueous				
☑ C hydrated				
□ D saturated				
(ii) When the product in equation 1 is gently heated, another reaction occurs. Equation 2 represents this reaction.				
Equation 2 $CoCl_2.6H_2O(s) \rightarrow CoCl_2.2H_2O(s) + 4H_2O(g)$				
What do equations 1 and 2 suggest about the reactions?				
reversible	(1)			
(Total for Question 2 = 7	marks)			

3 The halogens are elements in Group 7 of the Periodic Table.

The halogens react with metals to form compounds called halides.

Table 1 shows information about some halogens and their halides.

Halogen	Appearance at room temperature	Halide	Melting point in °C
chlorine	green gas	lithium chloride	605
bromine	red-brown liquid	sodium bromide	747
iodine	grey solid	potassium iodide	681

Table 1

		120 YES 10 DE		127			
(a)	(i)	Predict the	physical	state	of fluorine	at room	temperature.

(1)

(ii) Predict how the colour of a tatine at room temperature compares with the colour of iodine.

(1)

du-her

(b) Each of the halides in table 1 was dissolved in water to form a solution.

A sample of each of the halogens was then added to some of the halide solutions.

Table 2 shows the results.

Markala.	Halogen added		
Halide	Chlorine	Bromine	lodine
lithium chloride	not done	no reaction	no reaction
sodium bromide	orange solution	not done	no reaction
potassium iodide	brown solution	brown solution	not done


Table 2

(i) Suggest why there is no reason to add chlorine to lithium chloride solution. (1)

3 (ii) Why was there no reaction when iodine was added to sodium bromide solution	on? (1)
12 less reactive	
(iii) Name the substance with the brown colour that formed when chlorine was added to potassium iodide solution.	
	(1)
to lodine	
(iv) The reaction between bromine and potassium iodide solution is a displacement reaction.	
What is the correct description of this reaction?	
	(1)
☐ A bromide displaces iodide	
B bromine displaces iodide	
bromide displaces iodine	
D bromine displaces iodine	
(v) Complete the chemical equation for the reaction between chlorine and potassium bromide solution.	(1)
Cl ₂ + 2KBr → 2KC1 + BC2	V-7
(Total for Question 3 = 7 ma	rks)

- 4 Ethene can be converted into many useful substances.
 - (a) Draw a dot and cross diagram to show the covalent bonding in a molecule of ethene. Only the outer electrons in each atom need to be shown.

(2)



- (b) Compound X is made from ethene and is used in cars to prevent the engine coolant from freezing in cold weather.
 - (i) Compound X contains 38.7% carbon, 9.7% hydrogen and 51.6% oxygen by mass.

Calculate the empirical formula of X.

$$\frac{38.7}{12} \frac{4.7}{16} \frac{51.6}{16}$$

Empirical formula CH2O

(ii) The relative formula mass (M₂) of X is 62

What is the molecular formula of X?

$$\frac{62}{31} = 2$$

(1)

Molecular formula (2H602

(Total for Question 4 = 6 marks)

5 A student was asked to compare the industrial processes used to extract aluminium and iron from their ores.	4
(a) (i) Name the main ore used as the source of iron.	(1)
(ii) Aluminium is extracted from purified aluminium oxide.	
What is the formula of aluminium oxide?	
	(1)
A1203	
(iii) One solid element is used in the extraction of both metals.	
Identify this element and state its purpose in the extraction of aluminium.	(2)
Element	
Purpose electroles	
(iv) One gaseous element takes part in a reaction needed in the extraction of iron. Identify this element and state its purpose in the extraction of iron.	(2)
Purpose rodue had	
Purpose produce hast . formation of reducing agent.	



	TO A ROLL WANTE
(b) The student wrote this statement: The extractions of aluminium and iron both involve reduction of	and oxidation reactions.
(i) What name is given to a reaction that involves both reduction a	and oxidation? (1)
Keaox	
(ii) Why does this equation represent a reduction reaction?	
$Al^{3+} + 3e^- \rightarrow Al$	
71 1 3C 7 71	(1)
A13+ gains of eletrons	
(iii) The equation for a reaction that occurs in some extractions of in	on is
$C + H_2O \rightarrow CO + H_2$	
Identify the substance oxidised in this reaction, giving a reason	for your choice. (2)
Substance oxidised Carbon	
Reason gans oxygen.	

5 (c) Both extractions occur at a high temperature.

Neither extraction uses a catalyst.

(i) What is meant by the term catalyst?

(2)

· chemical inchanged / lower activation

(ii) State one reason why cryolite is used in the extraction of aluminium.

(1)

· Solvent

· lowers operating temp.

- (d) Several equations can be written for the reactions occurring in the extractions.
 - (i) Write the chemical equation for the reaction between iron(III) oxide (Fe₂O₃) and carbon monoxide (CO).

(2)

Fezoz +300 -> 2Fe + 300,

(ii) This equation represents a reaction used to remove impurities in the extraction of iron.

State the type of reaction occurring in this equation.

(1)

neutralisation

(iii) Complete the table by giving the common name for calcium silicate.

(1)

Formula of compound	Chemical name	Common name
CaO	calcium oxide	quicklime
CaSiO ₃	calcium silicate	Slag

(Total for Question 5 = 17 marks)

(3)

(2)

6 The table shows the structures of six organic compounds, A to F.

H	B H C=C H	CH ₃ CH ₃ CH ₃ CH ₃ CH ₃
H H HCCH H H	H H H	F H

- (a) The letter of the compound in the table that is **not** shown as a displayed formula is ______(1)
- (b) (i) State what is meant by the term hydrocarbon, and give the letter of one compound in the table that is **not** a hydrocarbon.

Hydrocarbon carbais only hydrogen + carbon

Letter A

(ii) State what is meant by the term unsaturated, and give the letter of one compound in the table that is unsaturated.

Unsaturated contains a double bond

Letter S

(iii) State what is meant by the term isomers, and give the letters of two compounds in the table that are isomers of each other.

Isomers some indecular formula different structure

Letters and

(c) Some of the compounds in the table are members of the same homologous series.
 (i) One feature of a homologous series is that adjacent members have formulae that differ by CH₂

State two other features of members of the same homologous series.

(2)

Some

general

formula

2 similar

chemical

properties

(ii) Give the letters of two adjacent members of the same homologous series shown in the table.

(1)

Letters

D

and

(d) (i) Compound G has the molecular formula C₂H₄Br₂

It can be made from a compound in the table by a reaction that does **not** need UV light.

Draw the displayed formula of compound G.

(1)

(ii) Compound H reacts with bromine to form one of the compounds in the table. The reaction needs UV light.

Draw the displayed formula of compound H.

(1)

(Total for Question 6 = 14 marks)


7 A teacher asked her students to suggest some experiments that could be done using chemicals found in the home. One student planned an experiment to measure the temperature change when baking soda is added to vinegar.

She wrote this plan.

- pour 100 cm³ of vinegar into a polystyrene cup
- weigh out five separate 1 g portions of baking soda
- measure the temperature of the vinegar
- add 1 g of baking soda to the vinegar and stir
- record the new temperature
- add the other portions of baking soda, stirring and recording the temperature after each portion is added

The graph shows her results.

of mixture in °C



Mass of baking soda added in grams

(a) The student said that the reaction in her experiment was not complete.	
How does the graph support her statement?	
line not level	(1)
(b) The student used a polystyrene cup rather than a glass beaker.	
Why is it better to use a polystyrene cup?	
insulator	(1)
(NSCACO)	***************************************
(c) Vinegar contains ethanoic acid. Baking soda contains sodium hydrogencarbona	te.
The student found this equation for the reaction:	
CH₃COOH + NaHCO₃ → CH₃COONa + H₂O + CO₂	
(i) There is no colour change during this reaction.	
Suggest one observation, other than the change in temperature, that could be made during the reaction.	
f: .	(1)
fizz.ng	
(ii) The compound CH ₃ COOH is an acid and the compound CH ₃ COONa is a salt.	
The graph shows that the temperature goes down during the reaction.	
Use this information to state the two types of reaction occurring.	
	(2)
1 endollermic	
2 neutralisation	

(1)

 \rightarrow (d) (i) Complete the energy level diagram by showing the products of the reaction.

CH3COON + H20 + CO2

AH

CH3COOH + NaHCO3

(ii) Label the diagram to show the energy change, ΔH , for the reaction.

(1)

(e) The student repeated the experiment using the same method with a different sample of vinegar. She recorded these results.

Volume of vinegar = 100 cm³

Mass of baking soda = 5.0 g

Temperature at start = 18.7 °C

Temperature at end = 13.2 °C

(i) Calculate the heat energy change in this experiment using the expression

heat energy change = volume of vinegar \times 4.2 \times temperature change

- 100 x 4.2 x 5.5

Heat energy change = 2310 J

(ii) The student wanted to calculate the amount, in moles, of ethanoic acid in the vinegar.

Apart from the volume of vinegar, what other information would she need to be able to calculate the amount of ethanoic acid?

(1)

(2)

concentration.

(Total for Question 7 = 10 marks)

BLANK PAGE

8 A group of students investigated the reaction between marble chips (calcium carbonate) and dilute hydrochloric acid.

The equation for this reaction is


$$CaCO_3(s) + 2HCI(aq) \rightarrow CaCI_2(aq) + H_2O(I) + CO_2(g)$$

They wanted to find the effect of changing the concentration of hydrochloric acid on the rate of reaction. The teacher provided a solution that she had labelled 100% hydrochloric acid.

The teacher told them to do all their experiments

- using different concentrations of hydrochloric acid made by diluting the 100% hydrochloric acid
- by timing how long it took to collect carbon dioxide
- at room temperature

The students used this apparatus.



(a) The students tried to keep the amount of calcium carbonate constant by using the same number of marble chips in each experiment.

State two other properties of the marble chips that should be the same in each experiment.

(2)

Temp Muss

2 surfue area



 \S (b) The table shows how some of the students wrote down their results.

Student	Results
1	I used 6 marble chips and 100% hydrochloric acid and collected 100 cm³ of gas in 40 seconds.
2	In my experiment there were 6 marble chips and 80% hydrochloric acid and I collected 100 cm³ of carbon dioxide by the end of the experiment.
3	The marble chips and 60% hydrochloric acid formed 100 cm³ of gas in 70 seconds.
4	I used 40% hydrochloric acid and 6 marble chips. It took 105 seconds to collect the gas.
5	I collected 100 cm³ of gas in 135 seconds when I used 6 marble chips.

The teacher said that she could only use the results from student 1 because the other students had not recorded enough information.

Identify the piece of information that each student failed to record.

Student 2 Time

Student 3 No- of Marble imps

Student 4 Vol of gas

Student 5 Cone of HC1


 $\hat{\xi}$ (c) The students repeated the experiment to make sure that it was a fair test. They used a different supply of hydrochloric acid. They all measured the time to collect 100 cm³ of carbon dioxide and calculated the rate of each reaction.


Their results are shown in the table.

Percentage concentration of hydrochloric acid	Time to collect 100 cm³ of gas in seconds	Rate of reaction in cm ³ /s
20	66.7	1.5
40	52.6	1.9
60	34.5	2.9
70	30.3	3.3
80	25.6	3.9
100	20.8	4.8

(i) Plot these results on the grid and draw a straight line of best fit.

(3)






(ii) One of the points is anomalous. Circle this point on the graph.

(1)

(iii) Suggest two errors in the experiment that could have caused this result.	s anomalous
. Timer starter lake	(2)
1	
o conc bo high	
. more marke chips.	
· air whendy in cylinder	
(iv) Use your graph to estimate the rate of reaction using an acid con Show on your graph how you obtained your answer.	centration of 50%.
Show on your graph now you obtained your answer.	(2)
2·4 cm³/s	(2)
(Total for Questic	on 8 = 14 marks)

9 The diagram shows some pieces of apparatus used to measure volumes.



A student was given a large bottle containing sodium hydroxide solution and a supply of dilute sulfuric acid of known concentration.

He was allowed to use normal laboratory apparatus, including the pieces of apparatus shown in the diagram.

He was told to plan an experiment to find the concentration of the sodium hydroxide solution.

This is his plan.

- Step 1 Obtain about 150 cm³ of each solution.
- Step 2 Use a measuring cylinder to add exactly 25.0 cm³ of sodium hydroxide solution to a conical flask.
- Step 3 Add a few drops of universal indicator to the conical flask.
- Step 4 Use a burette to add the sulfuric acid to the conical flask until the indicator changes colour.
- (a) (i) Give the name of the most suitable piece of apparatus in the diagram that should be used in Step 1.

(1)

bruker

(ii) Give the name of the piece of apparatus in the diagram that should be used instead of a measuring cylinder in Step 2.

(1)

pipette

(iii) State why universal indicator is **not** a good choice for this experiment and suggest an indicator that would be more suitable.

(2)

end point not shorp.

Methyl orange


(iv) Why is a pipette not suitable for adding the acid in Step 4?

(1)

counst be conholled

(b) The diagram shows the burette readings in one experiment before and after adding the acid.





Use the readings to complete the table, entering all values to the nearest 0.05 cm³.

(3)

Burette reading after adding acid in cm ³	2.75
Burette reading before adding acid in cm ³	22.60
Volume of acid added in cm³	19.85

(c) The student repeated the experiment using a different concentration of sodium hydroxide solution and recorded these results.

Burette reading after adding acid in cm ³	24.90	25.85	24.85	25.55
Burette reading before adding acid in cm ³	1.20	2.75	1.50	2.10
Volume of acid added in cm ³	23.70	23.10	23.35	23.45
Titration results to be used (✓)			~	V

The volumes of acid added during these titrations are not all the same. The average (mean) volume of acid should be calculated using only concordant results.

Concordant results are those volumes that differ from each other by 0.20 cm³ or less.

- (i) Identify the concordant results by placing ticks (✓) in the table where appropriate. (1)
- (ii) Use your ticked results to calculate the average (mean) volume of acid added.

Average (mean) volume of acid = $\frac{23.40}{\text{cm}^3}$ cm³

(d) The student used the same method to find the concentration of a solution of potassium hydroxide. The equation for the reaction is

$$2KOH + H_2SO_4 \rightarrow K_2SO_4 + 2H_2O$$

These are his results.

Volume of potassium hydroxide solution	25.0 cm ³
Volume of sulfuric acid	23.60 cm ³
Concentration of sulfuric acid	0.0500 mol/dm³

He used these results to calculate the concentration of the potassium hydroxide solution.

Step 1 amount of
$$H_2SO_4 = \frac{0.0500 \times 23.60}{100} = 0.0118 \,\text{mol}$$

Step 2 amount of KOH =
$$\frac{0.0118}{2}$$
 = 0.00590 mol

Step 3 concentration of KOH =
$$\frac{0.00590}{23.60} \times 1000 = 0.250 \text{ mol/dm}^3$$

There is one mistake in each step of the calculation.

What correction should the student make in each step?

(i) Step 1

(ii) Step 2

(1)

X2

(iii) Step 3

(1)

-25.0

(Total for Question 9 = 14 marks)

10 The Haber process is used to convert nitrogen and hydrogen into ammonia.			
(a) Which raw material is the source of the nitrogen?			
	(1)		
air	***************************************		
(b) The hydrogen for this process is obtained using reactions 1 and 2			
Reaction 1 $CH_4(g) + H_2O(g) \rightleftharpoons CO(g) + 3H_2(g)$ $\Delta H = +210 \text{ kJ mol}^{-1}$			
Reaction 2 $CO(g) + H_2O(g) \rightleftharpoons CO_2(g) + H_2(g)$ $\Delta H = -42 \text{ kJ mol}^{-1}$			
(i) Predict what will happen to the equilibrium position in reaction 1 when the pressure is increased. Give a reason for your prediction.	(2)		
Prediction Move to left	(2)		
Prediction Move 10 4/1			
Reason fewer molecules	an		
(ii) Predict what will happen to the equilibrium position in reaction 2 when the			
temperature is increased. Give a reason for your prediction.	(2)		
Prediction more to left			
Reason exotlermic reaction			
Reason Exoremic Teacher			
(c) Predict what will happen to the rate of reaction 1 when the pressure and temperature are increased.			
temperature are increased.	(2)		
Effect of increased pressure increased			
Effect of increased temperature in weal			

\□ (d) Some of the ammonia from the Haber process is made into fertilisers such as ammonium nitrate.		
The equation for the formation of ammonium nitrate is		
$NH_3 + HNO_3 \rightarrow NH_4NO_3$		
A manufacturer makes a batch of 40 kg of ammonium nitrate.		
(i) Calculate the amount, in moles, of NH ₄ NO ₃ in 40 kg of ammonium nitrate.		
	(2)	
40,000		
80/		
Amount of NH ₄ NO ₃		mol
(ii) Deduce the amount, in moles, of ammonia needed to make this amount of		11101
ammonium nitrate.	-Mark-Marca III	
	(1)	
Amount of ammonia Soo	***************************************	mol
(iii) Calculate the mass, in kg, of ammonia needed.	(2)	
0 - 0 - 0 - 0 - 0	V=1	
SOOX 1) = 8500		
Mass of ammonia 8.5		ka
(e) The manufacturer also produces fertilisers that are labelled NPK.		
Suggest the names of two elements, other than nitrogen, that are likely to be pre in NPK fertilisers.	esent	
	(1)	
Potassium and Phosophorons	***********************	
(Total for Question 10 = 13 m		


11	The properties of substances can be explained in terms of their bonding and structure.		
	Electricity can be transmitted by overhead power lines. This method of transmission requires electrical conductors and insulators.		
	(a) Aluminium is used for the overhead lines because it is a good conductor of electricity and is ductile (can be pulled into a wire).This diagram can be used to represent the structure of aluminium.		
	aluminium atom		
	(i) Explain why aluminium is a good conductor of electricity.		
	delocalised electrons that move		
	(ii) Suggest why aluminium is ductile. (2) layers of atoms that can slide (over each ofter)		

(5)


() The main substance that acts as an insulator in this method of transmission of electricity is air, which is mostly nitrogen.

The power lines are supported by solid insulators. Most solid insulators are manufactured using silica.

The diagram shows the structures of nitrogen and silica.



Nitrogen



Silica

Explain, in terms of bonding and structure, why nitrogen is a gas at room temperature but silica is a solid with a high melting point.

Nitrogen . (Simple) molecular smuchue
· weak internolecular forces (of athachins between
(or read little energy molecule).
to over come)
Silica giant covalent shuchues with
"Shong bonds or (road a look of enery to
breach).
(Total for Question 11 = 9 marks)
TOTAL FOR PAPER - 120 MARKS

BLANK PAGE