

Please check the examination details below before entering your candidate information

Candidate surname		Other names	
Centre Number		Candidate Number	
Pearson Edexcel International GCSE (9–1)		<div style="display: flex; justify-content: space-around;"> <div style="border: 1px solid black; width: 20px; height: 20px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px;"></div> </div> <div style="display: flex; justify-content: space-around;"> <div style="border: 1px solid black; width: 20px; height: 20px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px;"></div> </div>	
Thursday 9 January 2020			
Morning (Time: 2 hours)		Paper Reference 4CH1/1C 4SD0/1C	
Chemistry Unit: 4CH1 Science (Double Award) 4SD0 Paper: 1C			
You must have: Calculator, ruler			Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- Show all the steps in any calculations and state the units.
- Some questions must be answered with a cross in a box ☒. If you change your mind about an answer, put a line through the box ☒ and then mark your new answer with a cross ☒.

Information

- The total mark for this paper is 110.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P59923A

©2020 Pearson Education Ltd.

1/1/1/1/1/



The Periodic Table of the Elements

1	2	<div>1<div>1</div>H<div>1</div>hydrogen</div>										3	4	5	6	7	0
7	9	Key <div>relative atomic mass atomic symbol name atomic (proton) number</div>										11	12	14	16	19	20
Li lithium 3	Be beryllium 4											B boron 5	C carbon 6	N nitrogen 7	O oxygen 8	F fluorine 9	Ne neon 10
23	24											Al aluminium 13	Si silicon 14	P phosphorus 15	S sulfur 16	Cl chlorine 17	Ar argon 18
39	40											Ga gallium 31	Ge germanium 32	As arsenic 33	Se selenium 34	Br bromine 35	Kr krypton 36
85	88											In indium 49	Sn tin 50	Sb antimony 51	Te tellurium 52	I iodine 53	Xe xenon 54
133	137											Tl thallium 81	Pb lead 82	Bi bismuth 83	Po polonium 84	At astatine 85	[222] Rn radon 86
[223] Fr francium 87	[226] Ra radium 88											Elements with atomic numbers 112-116 have been reported but not fully authenticated					

* The lanthanoids (atomic numbers 58-71) and the actinoids (atomic numbers 90-103) have been omitted.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Answer ALL questions.

1 This question is about gases in the atmosphere.

(a) The box gives the names of some gases in the atmosphere.

argon	carbon dioxide	helium	nitrogen	oxygen
-------	----------------	--------	----------	--------

Choose gases from the box to answer these questions.

Each gas may be used once, more than once or not at all.

(i) Identify a noble gas.

(1)

Argon / Helium

(ii) Identify a gas that makes up about 78% of the atmosphere.

(1)

Nitrogen

(iii) Identify a greenhouse gas.

(1)

Carbon Dioxide

(iv) Identify a gas produced by the thermal decomposition of calcium carbonate.

(1)

Carbon Dioxide

(b) Sulfur reacts with oxygen to produce sulfur dioxide gas.

(i) Write a chemical equation for this reaction.

(1)



(ii) State an environmental problem caused when sulfur dioxide gas dissolves in water in the atmosphere.


(1)

Acid Rain.

(Total for Question 1 = 6 marks)



2 The diagram shows the electronic configuration of an atom of an element.



(a) Complete the table by giving the missing information about this atom.

(5)

name of the part of this atom labelled Z	Nucleus
number of protons in this atom	12
number of the group that contains this element	2
number of the period that contains this element	3
the charge on the ion formed from this atom	2+



(b) This element has three isotopes.

The table shows the mass number and percentage abundance of each isotope in a sample of this element.

Mass number	Percentage abundance (%)
24	79.2
25	10.0
26	10.8

Calculate the relative atomic mass (A_r) of this element.

Give your answer to one decimal place.


(3)

$$A_r = 24(0.792) + 25(0.1) + 26(0.108) \\ = 24.316$$

relative atomic mass = 24.3
(1.d.p.)

(Total for Question 2 = 8 marks)

3 The diagram shows two samples of iron, A and B.



Sample B is coated with a thin layer of zinc.

(a) Name the process used to coat iron with zinc.

(1)

Galvanising.

(b) The two samples of iron are left outside for several weeks.

A brown solid containing hydrated iron(III) oxide forms on sample A.

(i) Give the common name for the brown solid.

(1)

Rust

(ii) Give the names of the two substances that react with the iron to form the brown solid.

(2)

1

Water


2

Oxygen.



(c) Iron can be formed by reacting aluminium powder with iron(III) oxide.

The diagram shows how this reaction can be demonstrated.



When the magnesium fuse is lit, a very exothermic reaction occurs.

(i) State the meaning of the term **exothermic**.

(1)

A reaction which releases thermal energy.

(ii) The equation for the reaction between aluminium and iron(III) oxide is



Explain what this reaction shows about the relative reactivities of aluminium and iron.

(2)

Aluminium is more reactive than iron as aluminium displaces iron.

(iii) Explain why the reaction between aluminium and iron(III) oxide is a redox reaction.

(3)

Aluminium gains oxygen so is oxidised.

Iron loses oxygen so it is reduced.

Both oxidation and reduction occur, hence Redox.

(Total for Question 3 = 10 marks)



4 This question is about ionic compounds.

- (a) The table shows the formulae of some positive and negative ions, and the formulae of some compounds containing these ions.

	Mg^{2+}	Al^{3+}	NH_4^+
S^{2-}	MgS	Al_2S_3	$(\text{NH}_4)_2\text{S}$
NO_3^-	$\text{Mg}(\text{NO}_3)_2$	$\text{Al}(\text{NO}_3)_3$	NH_4NO_3
CO_3^{2-}	MgCO_3	$\text{Al}_2(\text{CO}_3)_3$	$(\text{NH}_4)_2\text{CO}_3$

- (i) Complete the table by giving the three missing formulae.

(3)

- (ii) Give the name of the compound with the formula NH_4NO_3

(1)

Ammonium nitrate.

- (b) Sodium oxide, Na_2O , is an ionic compound.

The sodium and oxide ions are held together by ionic bonds.

- (i) State the meaning of the term **ionic bond**.


(2)

The strong electrostatic attraction between oppositely charged ions.




- (ii) The diagram shows the arrangement of the electrons in a sodium atom and in an oxygen atom.

sodium atom




oxygen atom



Draw diagrams in the boxes to show the arrangement of the electrons in the ions of sodium oxide.







Include the charges on the ions.

(3)



(Total for Question 4 = 9 marks)

5 The boxes show the displayed formulae of six organic compounds, P, Q, R, S, T and U.

		
P	Q	R
		
S	T	U

(a) Use the letters P, Q, R, S, T and U to answer these questions.

Each letter may be used once, more than once or not at all.

(i) Give the letter of the compound that is **not** a hydrocarbon.

S

(1)

(ii) Give the letters of the two compounds that have the same empirical formula.

T, U

(1)

(iii) Give the letter of the compound that is used to manufacture poly(propene).

U

(1)



(b) Describe a test that can be used to distinguish between compounds Q and T.

(3)

test *Add bromine water.*

result with compound Q *No Colour Change*

result with compound T *Bromine decolourises from orange to Colourless.*

(c) Compounds P, Q and R are members of the same homologous series.

Give two characteristics of a homologous series.

(2)

1 *All members have the same general formula.*


Each successive member differs by CH_2 .

2 *Each member has the same functional group.*

Similar chemical properties of each member.

Trend in physical properties between successive members.

(d) This is the displayed formula of an alkene, V.




(i) Give the name of alkene V.

(1)

but-1-ene.

(ii) Draw the displayed formula of another alkene that is an isomer of alkene V.

(1)



(e) An organic compound has the percentage composition by mass


C = 36.36%

H = 6.06%

F = 57.58%

(i) Show that the empirical formula of the compound is CH_2F


(2)



(ii) The relative molecular mass (M_r) of the compound is 66.

Determine the molecular formula of the compound.

(2)



molecular formula = $\text{C}_2\text{H}_4\text{F}_2$


(Total for Question 5 = 14 marks)





6 A student uses this method to investigate the reaction of dilute hydrochloric acid with zinc.

- pour some dilute hydrochloric acid into a glass beaker
- record the initial temperature of the acid
- add a piece of zinc and stir the mixture
- record the temperature of the mixture after one minute

(a) Write a word equation for the reaction of dilute hydrochloric acid with zinc.



(b) The diagram shows the thermometer readings for this reaction.



Complete the table, giving all values to the nearest 0.1 °C.

(3)

temperature in °C after adding zinc	22.4
temperature in °C before adding zinc	17.7
temperature change in °C	4.7

(c) Another student repeats the method using five different metals to compare their reactivity.

(i) This student uses a polystyrene cup instead of a glass beaker.

Explain why a polystyrene cup is better than a glass beaker in this investigation.

(2)

Polystyrene is an insulator so reduces heat loss.

Hence, the temperature change will be closer to the true value.

(ii) Give three factors that the student should keep constant in this investigation.

(3)

1 Amount of metal

2 Concentration of acid.

3 Volume of acid.

Room temperature

Speed and time of stirring.



(d) The table shows some of the student's results.

Metal added	Observation	Temperature change in °C
copper	no bubbling	0.0
iron	slow bubbling	
magnesium	rapid bubbling	8.7
tin	very slow bubbling	1.4
zinc	moderate bubbling	5.1

(i) State why there is no temperature change for copper.

(1)

No reaction occurred.

(ii) Predict the temperature change for iron.

(1)

(1.5 to 5.0 °C)

temperature change = 2.0 °C

(iii) Deduce the order of reactivity of the five metals.

(1)

most reactive Magnesium

Zinc

Iron

Tin

least reactive Copper

(Total for Question 6 = 12 marks)

7 This question is about some of the halogens and their compounds.

(a) (i) Which element is a liquid at room temperature?

(1)

- ☐ A astatine
- ☒ B bromine
- ☐ C chlorine
- ☐ D iodine

(ii) Which element has the palest colour?

(1)

- ☐ A astatine
- ☐ B bromine
- ☒ C chlorine
- ☐ D iodine

(iii) Which element is the least reactive?

(1)

- ☒ A astatine
- ☐ B bromine
- ☐ C chlorine
- ☐ D iodine



(b) A teacher uses displacement reactions to demonstrate the reactivities of some halogens.

She adds solutions of chlorine, bromine and iodine separately to three different sodium halide solutions.

The table shows some of the teacher's results.

	sodium chloride	sodium bromide	sodium iodide
chlorine solution	not done	solution turns orange	<i>Solution turns brown.</i>
bromine solution	solution stays orange	not done	solution turns brown
iodine solution	<i>Solution stays brown</i>	solution stays brown	not done

A change in colour of the solution indicates that a reaction has occurred.

(i) Complete the table by predicting the missing results.

(2)

(ii) State why the teacher does not add bromine solution to sodium bromide solution.

(1)

Bromine cannot displace itself.

(iii) The word equation for the reaction of bromine with sodium iodide is

bromine + sodium iodide → iodine + sodium bromide

Write a chemical equation for this reaction.

(1)



(c) A technician sees an unlabelled bottle containing a liquid.

He knows that the liquid is a solution of one of these compounds.

- copper(II) chloride
- copper(II) bromide
- iron(II) chloride
- iron(II) bromide

Describe chemical tests that the technician could use to identify the compound in the solution.

Add Sodium hydroxide Solution to a Sample of the liquid. ⁽⁶⁾ If a blue precipitate forms, Cu^{2+} ions are present. If a green precipitate forms, Fe^{2+} ions are present.

Add Silver nitrate Solution to a Sample of the liquid. If a white precipitate forms, Chloride ions are present. If a Cream precipitate forms, bromide ions are present.

(Total for Question 7 = 13 marks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



8 (a) Carbon dioxide changes directly from a solid to a gas without becoming a liquid.

(i) Give the name of the change of state from solid to gas.

(1)

Sublimation.

(ii) Describe the test for carbon dioxide gas.

(2)

~~Add~~ Bubble the gas into limewater. If CO_2 (g) is present, the limewater turns cloudy.

(b) Carbon dioxide is a simple molecular covalent substance.

Explain why carbon dioxide turns from a solid to a gas at a very low temperature.

(2)

Weak intermolecular forces require little energy to overcome, achieved by low temperatures.



(c) Diamond and graphite are both giant covalent substances made up of carbon atoms.

- diamonds are used in cutting tools
- graphite is used in pencils to make marks on paper

Explain, with reference to structure and bonding, why each substance is suitable for its particular use.

(6)

In diamond, each Carbon atom is Covalently bonded to four other Carbon atoms in a giant tetrahedral lattice structure. The covalent bonds are very strong, making diamond very hard & suitable for cutting tools.

In Graphite, each Carbon atom is Covalently bonded to 3 other Carbon atoms. The structure is in layers with weak forces of attraction between the layers so the layers can slide over each other, making graphite soft - suitable for making marks on paper.

(Total for Question 8 = 11 marks)

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



- 9 A student uses this apparatus to investigate the heat energy released when a liquid fuel is burned.



This is the student's method.

- measure the mass of the spirit burner and fuel
- add 100 cm^3 of water to the copper can
- record the temperature of the water
- use the spirit burner to heat the water until the temperature rises by 30°C
- immediately measure the new mass of the spirit burner and fuel

- (a) Suggest why the student measures the mass of the spirit burner and fuel immediately after heating the water.

(1)

This reduces the mass loss due to evaporation of the liquid fuel.

- (b) When the fuel is burned, the student notices that a black solid forms on the bottom of the copper can.

- (i) Identify the black solid.

(1)

Carbon (Soot)

- (ii) Explain why the black solid forms.

(2)

Incomplete Combustion occurs as there is a limited oxygen supply.

- (c) (i) Show that the heat energy change, Q , to raise the temperature of 100 cm^3 of water by 30°C is approximately 13 kJ .

[mass of 1.0 cm^3 of water = 1.0 g]

[c for water = $4.2\text{ J/g/}^\circ\text{C}$]

(3)

$$\begin{aligned} Q &= mc\Delta T \\ &= 100(30)(4.2) \\ &= 12600\text{ J} \\ &= 12.6\text{ kJ} \\ &= \underline{13\text{ kJ}} \quad (2\text{ sig fig}) \end{aligned}$$

- (ii) The student burns 0.96 g of methanol, CH_3OH

Calculate the molar enthalpy change, ΔH , in kJ/mol , for the combustion of methanol.

Include a sign in your answer.

[M_r of methanol = 32]

(3)

$$n(\text{CH}_3\text{OH}) = \frac{\text{mass}}{M_r} = \frac{0.96}{32} = 0.03$$

$$\Delta H = \frac{-13}{0.03} = -433.333\text{ kJ mol}^{-1}$$

$$\Delta H = \underline{-433} \text{ kJ/mol} \quad (3\text{ sig fig})$$

- (d) The table shows data book values for the molar enthalpy change, ΔH , for the combustion of some alcohols with different numbers of carbon atoms per molecule.


Number of carbon atoms per molecule	1	2	3	4	5
Molar enthalpy change, ΔH , in kJ/mol	-730	-1370	-2020	-2680	-3320



- (i) Plot the data values from the table on the grid.

Draw a straight line of best fit.

(2)



- (ii) Deduce the value of ΔH for an alcohol with six carbon atoms per molecule.

Show on the graph how you obtained your answer.

(2)

$\Delta H = -4000$ kJ/mol

- (iii) State the relationship between ΔH and the number of carbon atoms per molecule.

(1)

The greater the number of carbon atoms per molecule, the more exothermic the value of ΔH is.

(Total for Question 9 = 15 marks)

10 Nitric acid (HNO_3) is used in the production of fertilisers.

Nitric acid is manufactured in three stages.

Stage 1 ammonia reacts with oxygen in the presence of a platinum catalyst to produce nitrogen monoxide gas, NO , and water.

Stage 2 nitrogen monoxide gas reacts with more oxygen to produce nitrogen dioxide gas, NO_2 .

Stage 3 nitrogen dioxide gas reacts with water to produce nitric acid and more nitrogen monoxide gas.

(a) (i) Complete the chemical equation for the reaction in stage 1.



(ii) Give the meaning of the symbol \rightleftharpoons

Reversible reaction. (1)

(iii) State the purpose of the platinum catalyst.

Increase the rate of reaction. (1)

(b) Give a chemical equation for the reaction of nitrogen monoxide and oxygen in stage 2.



(c) (i) The equation for the reaction in stage 3 is



Calculate the maximum mass, in tonnes, of nitric acid that could be produced in this reaction from 11.5 tonnes of nitrogen dioxide.

[1 tonne = 1.0×10^6 g]

(4)

$$n(\text{NO}_2) = \frac{\text{mass}}{M_r} = \frac{11.5 \times 10^6}{14 + 2(16)} = \frac{250\,000}{3} \text{ mol}$$

$$\text{Ratio } n(\text{NO}_2 : \text{HNO}_3) = 3 : 2$$

$$\therefore n(\text{HNO}_3) = \frac{250\,000}{3} \times 2$$

$$= \frac{500\,000}{3} \text{ mol}$$

$$M(\text{HNO}_3) = M_r \times n$$
$$= \frac{500\,000}{3} \times (1 + 16 + 3(16)) = 10.5 \times 10^6 \text{ g}$$

mass of nitric acid = 10.5 tonnes

(ii) Suggest what use can be made of the nitrogen monoxide gas formed in stage 3.


(1)

It can be reused in stage 2 to make more nitrogen dioxide.

QUESTION 10 CONTINUES ON NEXT PAGE

(d) When copper(II) oxide reacts with dilute nitric acid, copper(II) nitrate is produced.

The equation for the reaction is



0.200 mol of nitric acid reacts with excess copper(II) oxide.

A mass of 15.3 g of copper(II) nitrate is produced.

Calculate the percentage yield of copper(II) nitrate.

[M_r of copper(II) nitrate = 187.5]

(3)

$$n(\text{HNO}_3) = 0.2 \text{ mol}$$

$$n(\text{Cu}(\text{NO}_3)_2) = \frac{0.2}{2} = 0.1$$

$$m(\text{Cu}(\text{NO}_3)_2) = M_r \times n = 187.5 \times 0.1 \\ = 18.75 \text{ g}$$

$$\% \text{ yield} = \frac{15.3}{18.75} \times 100 = 81.6\%$$

percentage yield = 81.6 %

(Total for Question 10 = 12 marks)

TOTAL FOR PAPER = 110 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

